arXiv:hep-th/0411149v2 26 Dec 2004

`hep-th/0411149`
TIT/HEP-532

# Conformal Sigma Models with Anomalous Dimensions and Ricci Solitons

**Muneto Nitta**[*]

*Department of Physics, Tokyo Institute of Technology*
*Tokyo 152-8551, JAPAN*

## Abstract

We present new non-Ricci-flat Kähler metrics with $U(N)$ and $O(N)$ isometries as target manifolds of superconformally invariant sigma models with an anomalous dimension. They are so-called Ricci solitons, special solutions to a Ricci-flow equation. These metrics explicitly contain the anomalous dimension and reduce to Ricci-flat Kähler metrics on the canonical line bundles over certain coset spaces in the limit of vanishing anomalous dimension.

---

[*] E-mail: `nitta@th.phys.titech.ac.jp`

# 1 Introduction

String compactification with consistent non-trivial background is one of important subjects for long time. Sigma model approach gives a set of equations of motion for such backgrounds by vanishing condition on the beta function [1]. Calabi-Yau manifolds, Ricci-flat Kähler manifolds, are required as compactified manifolds with constant dilaton backgrounds. However no explicit metrics for compact Calabi-Yau manifold are known. On the other hand, if we allow non-compact manifolds, some explicit metrics for non-compact Calabi-Yau manifolds can be constructed [2]–[11]. Explicit solutions for non-Ricci-flat Kähler manifold with a non-trivial dilaton background were obtained by Kiritsis, Kounnas and Lust [12] as a generalization of two dimensional Euclidean black hole [13, 14].

We can consider nonlinear sigma models whose scalar fields have an anomalous dimension instead of a constant dilaton background, because they can be transformed to each other by a field redefinition for the scalar fields to absorb the dilaton field [14]. Using the Wilsonian renormalization group equation, the beta function of the $\mathcal{N} = (2, 2)$ supersymmetric nonlinear sigma models consisting of the complex scalar fields $(\varphi^i, \varphi^{*i})$ with an anomalous dimension $\gamma$ was obtained in [15, 16, 17] (see [18] for a review) as[1]

$$-\frac{\partial g_{ij^*}}{\partial t} = \beta(g_{ij^*}) = \frac{1}{2\pi} R_{ij^*} + \gamma\Big(\varphi^k g_{lj^*}\Gamma^l{}_{ik} + \varphi^{*k} g_{il^*}\Gamma^{l^*}{}_{j^*k^*} + 2g_{ij^*}\Big) \tag{1.1}$$

with $g_{ij^*}$, $R_{ij^*}$ and $\Gamma^i{}_{ij}$ being the Kähler metric, the Ricci-form and the connection, respectively. Conformally invariant models are defined by the condition of vanishing beta function

$$R_{ij^*} + 2\pi\gamma\Big(\varphi^k g_{lj^*}\Gamma^l{}_{ik} + \varphi^{*k} g_{il^*}\Gamma^{l^*}{}_{j^*k^*} + 2g_{ij^*}\Big) = 0. \tag{1.2}$$

A $U(N)$-invariant solution for this equation was obtained in [16]. It was shown to be equivalent to the $U(N)$-invariant solution with a dilaton background constructed in [12]. This is due to the equivalence of this model to the dilaton model. In this note we derive another $U(N)$-invariant solution and its extension to $O(N)$.

Eq. (1.1) is a so-called Ricci-flow equation which has attracted much attention recently in mathematics (see [20] for a review by a physicist). In the Riemann manifold with a metric $g_{\mu\nu}$ and some vector field $\xi^\mu$, the general Ricci-flow equation is written as

$$-\frac{\partial g_{\mu\nu}}{\partial t} = R_{\mu\nu} + \nabla_\mu\xi_\nu + \nabla_\nu\xi_\mu. \tag{1.3}$$

If we take $\xi^\mu = 2\pi\gamma(\varphi^i, \varphi^{*i})$ as a special case in the Kähler manifold, the Ricci-flow equation (1.3) reduces to Eq. (1.1). Solutions to $\frac{\partial g_{\mu\nu}}{\partial t} = R_{\mu\nu} + \nabla_\mu\xi_\nu + \nabla_\nu\xi_\mu = 0$ are called Ricci solitons and play a central role in classification of manifolds. Our new solutions presented in this note are Kähler Ricci solitons and we hope that these solutions are useful for classification of Kähler manifolds.

---

[1] To derive this equation, it was useful to expand the Lagrangian in terms of Kähler normal coordinates [19] which are natural extension of Riemann normal coordinates to Kähler manifolds.

## 2 $U(N)$ Invariant Model

We prepare an $N$-vector $\vec{\phi} = (\phi^1, \cdots, \phi^N)$ belonging to the fundamental representation of $U(N)$. Let us assume that the Kähler potential $K$ is written as a function of the $U(N)$-invariant as

$$K = K(X)\,, \qquad X \equiv \vec{\phi}^\dagger \vec{\phi}. \tag{2.1}$$

Geometric quantities can be calculated [16], to yield

$$\begin{aligned}
g_{ij^*} &= K'\delta_{ij} + K''\phi^{*i}\phi^j && (2.2)\\
R_{ij^*} &= -\partial_i\partial_{j^*}\mathrm{tr}\,\log g_{kl^*}\\
&= -\left[(N-1)\frac{K''}{K'} + \frac{2K''+K'''X}{K'+K''X}\right]\delta_{ij}\\
&\quad -\left[(N-1)\left(\frac{K'''}{K''} - \frac{(K'')^2}{(K')^2}\right) + \frac{3K'''+K''''X}{K'+K''X} - \frac{(2K''+K'''X)^2}{(K'+K''X)^2}\right]\phi^{*i}\phi^j, && (2.3)\\
g_{lj^*}\Gamma^l{}_{ik} &= g_{ij^*,k} = K''(\phi^{*k}\delta_{ij} + \phi^{*i}\delta_{kj}) + K'''\phi^{*k}\phi^{*i}\phi^j && (2.4)
\end{aligned}$$

with the prime denoting a differentiation with respect to $X$. Substituting these into Eq. (1.2), we obtain an ordinary differential equation, from a term proportional to $\delta_{ij}$,

$$(N-1)\frac{K''}{K'} + \frac{2K''+K'''X}{K'+K''X} + a(K'+K''X) = 0 \tag{2.5}$$

with $a$ a constant defined by the anomalous dimension as $a \equiv -4\pi\gamma$, and the derivative of this equation with respect to $X$ from a term proportional to $\phi^{*i}\phi^j$. The differential equation (2.5) can be integrated to give

$$(K')^{N-1}(K'+K''X) = c\,e^{-aK'X} \tag{2.6}$$

with $c$ an integration constant. Defining the function

$$F \equiv K'X, \tag{2.7}$$

Eq. (2.6) can be rewritten as

$$F^{N-1}F'X^{1-N} = c\,e^{-aF}. \tag{2.8}$$

Again this can be integrated to give the algebraic equation

$$e^{aF}\sum_{r=0}^{N-1}(-1)^r\frac{(N-1)!\,F^{(N-1)-r}}{(N-1-r)!\,a^{r+1}} = \frac{c}{N}X^N + b \tag{2.9}$$

with $b$ an integration constant. This reproduces the solution found in [12, 16] with a boundary condition $F(0) = 0$ which implies $K(0) = \text{const}$. It reduces to $F = \frac{1}{a}\log(1+aX)$ for $N = 1$ which defines the two dimensional Euclidean black hole [13, 14] for $a > 0$.

We now construct a new $U(N)$-invariant solution. To this end it is useful to define new coordinates by

$$\vec{\phi}^T = \sigma(1, \vec{z}^T). \tag{2.10}$$

We label $z$ by the same indices $i, j, \cdots$ with $\phi$ in the following. Then the invariant can be rewritten as

$$X = \vec{\phi}^\dagger \vec{\phi} = |\sigma|^2(1 + |\vec{z}|^2) \equiv |\sigma|^2 Z\,. \tag{2.11}$$

It is useful to write down the metric in these coordinates as

$$\begin{aligned}
g &= \begin{pmatrix} g_{\sigma\sigma} & g_{\sigma j^*} \\ g_{i\sigma^*} & g_{ij^*} \end{pmatrix}, \\
g_{\sigma\sigma} &= K'Z + K''|\sigma|^2 Z^2, \\
g_{\sigma j^*} &= K'\sigma^* \partial_{j^*} Z + K''\sigma^* |\sigma|^2 Z \partial_{j^*} Z, \\
g_{ij^*} &= K'|\sigma|^2 \partial_i \partial_{j^*} Z + K''|\sigma|^4 \partial_i Z \partial_{j^*} Z.
\end{aligned} \tag{2.12}$$

Using a solution $F$ of the same equation (2.9) with a boundary condition $F(0) = \text{const.} \neq 0$ *different* from the previous one, the metric can be calculated in these coordinates as

$$\begin{aligned}
ds^2 &= c\, e^{-aF} F^{1-N} Z^N |\sigma|^{2N-2} |d\sigma|^2 + [c e^{-aF} F^{1-N} Z^{N-1} |\sigma|^{2N-2} \sigma^* \partial_{j^*} Z d\sigma dz^{*j} + \text{c.c.}] \\
&\quad + [F(Z^{-1}\partial_i \partial_{j^*} Z - Z^{-2} \partial_i Z \partial_{j^*} Z) + c e^{-aF} F^{1-N} Z^{N-2} |\sigma|^{2N} \partial_i Z \partial_{j^*} Z] dz^i dz^{*j}.
\end{aligned} \tag{2.13}$$

Due to that boundary condition, this has a coordinate singularity in the limit $\sigma \to 0$. We perform the coordinate transformation

$$\rho = \sigma^N / N \tag{2.14}$$

to remove this coordinate singularity. We thus obtain the final form

$$\begin{aligned}
ds^2 &= c\, e^{-aF} F^{1-N} Z^N |d\rho|^2 + [c N e^{-aF} F^{1-N} Z^{N-1} \partial_{j^*} Z \rho^* d\rho dz^{*j} + \text{c.c.}] \\
&\quad + [F(Z^{-1}\partial_i \partial_{j^*} Z - Z^{-2} \partial_i Z \partial_{j^*} Z) + c N^2 e^{-aF} F^{1-N} Z^{N-2} |\rho|^2 \partial_i Z \partial_{j^*} Z] dz^i dz^{*j}
\end{aligned} \tag{2.15}$$

with $Z = 1 + |\vec{z}|^2$.

The metric at the $\rho = 0$ surface

$$ds^2|_{\rho=0} = F(0)(Z^{-1}\partial_i \partial_{j^*} Z - Z^{-2} \partial_i \partial_{j^*} Z) dz^i dz^{*j} = F(0) \partial_i \partial_{j^*} \log(1 + |z|^2) dz^i dz^{*j} \tag{2.16}$$

is the Fubini-Study metric on $\mathbf{C}P^{N-1} \simeq SU(N)/[SU(N-1) \times U(1)]$. Therefore the metric (2.15) is the (canonical) line bundle over $\mathbf{C}P^{N-1}$, $\mathbf{C} \ltimes \mathbf{C}P^{N-1}$. In fact, if we take $a = 0$ we recover the Ricci-flat Kähler metric of Calabi [2] (see also [8]).

In the limit of the boundary condition constant $F(0)$ tending to zero, $\mathbf{C}P^{N-1}$ with the metric (2.16) shrinks and the whole metric (2.15) contains a singularity. It is an orbifold singularity in the orbifold $\mathbf{C}^N/\mathbf{Z}_N$ defined by the identification (2.14). Therefore $F(0)$ is a blow-up parameter for the orbifold singularity.

Since the asymptotic form of $F$ is $F \simeq \frac{N}{a}\log aX$ for large $X$, the metric becomes asymptotically

$$\begin{aligned} ds^2 \simeq\ & \frac{c}{aX}(NX\log aX)^{1-N}Z^N|d\rho|^2 + \left[\frac{cN}{aX}(NX\log aX)^{1-N}Z^{N-1}\partial_{j^*}Z\rho^* d\rho dz^{*j} + \text{c.c.}\right] \\ & + \left[\frac{N}{a}\log aX(Z^{-1}\partial_i\partial_{j^*}Z - Z^{-2}\partial_i Z\partial_{j^*}Z) + cN^2\left(\frac{\log aX}{aX}\right)^{1-N} Z^{N-2}|\rho|^2\partial_i Z\partial_{j^*}Z\right] dz^i dz^{*j}. \end{aligned} \tag{2.17}$$

The difference between the previous solution [12, 16] and the present solution is locally just the boundary condition. For the previous case, they required regularity on $K$ at $X = 0$, $K(0) = \text{const.}$ (or $F(0) = 0$), and therefore $K = k_0 + k_1X + k_2X^2 + \cdots$. It is, however, not necessary for regularity on the metric as seen above. For our case the condition is $F(0) = \text{const.}$, and therefore $K = k_{-1}\log X + k_0 + k_1X + k_2X^2 + \cdots$ is not regular at $X \to 0$. As a result the topology is drastically changed. The previous solution has topology $\mathbf{C}^N$, but the present solution has topology $\mathbf{C} \ltimes \mathbf{C}P^{N-1}$ blowing up the orbifold singularity in $\mathbf{C}^N/\mathbf{Z}_N$.

## 3 $O(N)$ Invariant Model

Let us generalize the solution obtained in the last section to a $O(N)$-invariant solution. We prepare an $N$-vector $\vec{\phi} = (\phi^A, \cdots, \phi^N)$ again and put a constraint

$$\vec{\phi}^2 = \sum_{A=1}^{N}(\phi^A)^2 = 0 \tag{3.1}$$

to define a conifold [3]. It is convenient to rewrite this constraint as

$$\vec{\phi}^T J\vec{\phi} = 0, \quad J \equiv \begin{pmatrix} 0 & \mathbf{0} & 1 \\ \mathbf{0} & \mathbf{1}_{N-2} & \mathbf{0} \\ 1 & \mathbf{0} & 0 \end{pmatrix} \tag{3.2}$$

with $J$ the rank-2 $O(N)$ invariant tensor. The constraint (3.1) can be solved as [21, 22]

$$\vec{\phi}^T = \sigma\left(1, z^i, -\frac{1}{2}\vec{z}^2\right). \tag{3.3}$$

Ricci-flat metrics on conifolds with the singularity deformed by $\vec{\phi}^2 = r$ were constructed in [3] for $N = 3$ and [4] for general $N$. Ricci-flat metrics on conifolds without deformation was constructed in

[6, 8] which is still regular by an integration constant. Our solution here is a non-Ricci-flat deformation of the latter one.

The $O(N)$-invariant can be written as

$$X \equiv \vec{\phi}^\dagger \vec{\phi} = |\sigma|^2 \left( 1 + |\vec{z}|^2 + \frac{1}{4} |\vec{z}^2|^2 \right) \equiv |\sigma|^2 Z. \tag{3.4}$$

The expression of the metric is the same as the metric (2.12) but with $X$ and $Z$ in Eq. (3.4). Components of the Ricci-form are calculated as

$$\begin{aligned} R_{\sigma\sigma^*} &= -L^{-1} L' Z - L^{-2} [L L'' - (L')^2] |\sigma|^2 Z^2, \\ R_{\sigma j^*} &= -L^{-1} L' \sigma^* \partial_{j^*} Z - L^{-2} [L L'' - (L')^2] \sigma^* |\sigma|^2 Z \partial_{j^*} Z, \\ R_{i j^*} &= -L^{-1} L' |\sigma|^2 \partial_i \partial_{j^*} Z - L^{-2} [L L'' - (L')^2] |\sigma|^4 \partial_i Z \partial_{j^*} Z \end{aligned} \tag{3.5}$$

with $L \equiv (K')^{N-2} (K'' X^2 + K' X)$. Components of the connection are

$$\begin{aligned} g_{\sigma\sigma^*,\sigma^*} &= K'' 3\sigma Z^2 + K''' \sigma |\sigma|^2 Z^3, \\ g_{\sigma\sigma^*,i^*} &= K' \partial_{i^*} Z + K'' 3 |\sigma|^2 Z \partial_{i^*} Z + K''' |\sigma|^4 Z^2 \partial_{i^*} Z, \\ g_{ij^*,\sigma^*} &= K' \sigma \partial_i \partial_{j^*} Z + K'' (2\sigma |\sigma|^2 Z \partial_i \partial_{j^*} Z + \sigma |\sigma|^2 \partial_i Z \partial_{j^*} Z) + K''' \sigma |\sigma|^4 Z \partial_i Z \partial_{j^*} Z, \\ g_{ij^*,k^*} &= K' |\sigma|^2 \partial_i \partial_{j^*} \partial_{k^*} Z + K'' |\sigma|^4 (\partial_{k^*} Z \partial_i \partial_{j^*} Z + \partial_i Z \partial_{j^*} \partial_{k^*} Z + \partial_{j^*} Z \partial_i \partial_{k^*} Z) \\ &\quad + K''' |\sigma|^6 \partial_i Z \partial_{j^*} Z \partial_{k^*} Z. \end{aligned} \tag{3.6}$$

Substituting all these quantities to Eq. (1.2), we get

$$(\log L)' = -a (K' X)'. \tag{3.7}$$

By integrating this we find that $F = K'X$ satisfies the equation similar to the $U(N)$ case,

$$F^{N-2} F' X^{3-N} = c\, e^{-aF} \tag{3.8}$$

with an integration constant $c$. Again this can be integrated to yield

$$e^{aF} \sum_{r=0}^{N-2} (-1)^r \frac{(N-2)!\, F^{N-2-r}}{(N-2-r)!\, a^{r+1}} = \frac{c}{N-2} X^{N-2} + b \tag{3.9}$$

with an integration constant $b$. The coordinate transformation

$$\rho = \frac{\sigma^{N-2}}{N-2} \tag{3.10}$$

is needed to remove the coordinate singularity. We thus obtain the final form of the metric

$$\begin{aligned} ds^2 &= c\, e^{-aF} F^{2-N} Z^{N-2} |d\rho|^2 + [c(N-2) e^{-aF} F^{2-N} Z^{N-3} \partial_{j^*} Z \rho^* d\rho dz^{*j} + \text{c.c.}] \\ &\quad + [F(Z^{-1} \partial_i \partial_{j^*} Z - Z^{-2} \partial_i Z \partial_{j^*} Z) + c\,(N-2)^2 e^{-aF} F^{2-N} Z^{N-4} |\rho|^2 \partial_i Z \partial_{j^*} Z] dz^i dz^{*j} \end{aligned} \tag{3.11}$$

with $Z = 1 + |\vec{z}|^2 + \frac{1}{4}|\vec{z}^2|^2$.

The metric at the $\rho = 0$ surface is

$$ds^2|_{\rho=0} = F(0)(Z^{-1}\partial_i\partial_{j^*}Z - Z^{-2}\partial_i\partial_{j^*}Z)dz^i dz^{*j} = F(0)\partial_i\partial_{j^*}\log\left(1 + |z|^2 + \frac{1}{4}|\vec{z}^2|^2\right) dz^i dz^{*j}. \quad (3.12)$$

This is the metric on the quadric surface $Q^{N-2} \simeq SO(N)/[SO(N-2) \times U(1)]$ [21, 22]. The metric (3.11) is thus the (canonical) line bundle over $Q^{N-2}$. The Ricci-flat metric on a conifold [6, 8] is obtained by taking $a = 0$ in the metric (3.11). Our metric is non-Ricci-flat deformation of that conifold.

The asymptotic form of the metric (3.11) is ($F \simeq \frac{N-2}{a}\log aX$)

$$\begin{aligned} ds^2 \simeq\ & c[(N-2)X\log aX]^{2-N}Z^{N-2}|d\rho|^2 \\ & + \left[c(N-2)\{(N-2)X\log aX\}^{2-N}Z^{N-3}\partial_{j^*}Z\rho^* d\rho dz^{*j} + \text{c.c.}\right] \\ & + \Big[\frac{N-2}{a}\log aX(Z^{-1}\partial_i\partial_{j^*}Z - Z^{-2}\partial_i Z\partial_{j^*}Z) \\ & \qquad + c(N-2)^2\left\{(N-2)X\log aX\right\}^{2-N}Z^{N-4}|\rho|^2\partial_i Z\partial_{j^*}Z\Big]\, dz^i dz^{*j}. \end{aligned} \quad (3.13)$$

## 4 Conclusion

We have given two new metrics (2.15) and (3.11) with $U(N)$ and $O(N)$ symmetries, respectively as solutions of Eq. (1.2) for target spaces of conformally invariant sigma models with an anomalous dimension $\gamma$. They are canonical line bundles over the projective space $\mathbf{C}P^{N-1}$ and the quadric surface $Q^{N-2}$, respectively. These metrics contain the anomalous dimension explicitly through the parameter $a = -4\pi\gamma$. In the limit of vanishing anomalous dimension $a \to 0$, they reduce to those for Calabi-Yau manifolds [6, 8]. Generalization to other base coset spaces [8, 9] is straightforward. These new solutions give examples of Kähler Ricci solitons, singular solutions to the Ricci-flow equation (1.3).

A paper [23] recently posted to ArXiv has overlap with the present work. They have obtained explicit solutions with a dilaton background which also reduce to the same Calabi-Yau manifolds [6]–[10] with the present work. Therefore solutions in the present paper and those in [23] would be equivalent to each other by some field redefinition.

## Acknowledgements

The author would like to thank Kiyoshi Higashijima and Etsuko Itou for a collaboration in the early stages of this work and for explanation of their works [15, 16]. He is also grateful to Takashi Maeda for explanation of the Ricci-flow equation. His work is supported by Japan Society for the Promotion of Science under the Post-doctoral Research Program.

# References

[1] J. Polchinski, "String theory. Vol. 1: An introduction to the bosonic string"; "String theory. Vol. 2: Superstring theory and beyond".

[2] E. Calabi, Ann. Scient. Ec. Norm. Sup. **12**, 269 (1979).

[3] P. Candelas and X. C. de la Ossa, Nucl. Phys. B**342**, 246 (1990); R. Minasian and D. Tsimpis, Nucl. Phys. B**572**, 499 (2000), [arXiv:hep-th/9911042]; K. Ohta and T. Yokono, JHEP **0002** (2000) 023, [arXiv:hep-th/9912266].

[4] M. B. Stenzel, Manuscripta Math. **80**, 151 (1993); M. Cvetic, G. W. Gibbons, H. Lu and C. N. Pope, Commun. Math. Phys. **232**, 457 (2003) [arXiv:hep-th/0012011]; K. Higashijima, T. Kimura and M. Nitta, Phys. Lett. B **515**, 421 (2001) [arXiv:hep-th/0104184].

[5] D. N. Page and C. N. Pope, Class. Quantum Grav. **4** (1987) 213.

[6] K. Higashijima, T. Kimura and M. Nitta, Phys. Lett. B **518**, 301 (2001) [arXiv:hep-th/0107100].

[7] K. Higashijima, T. Kimura and M. Nitta, Nucl. Phys. B **623**, 133 (2002) [arXiv:hep-th/0108084].

[8] K. Higashijima, T. Kimura and M. Nitta, Annals Phys. **296**, 347 (2002) [arXiv:hep-th/0110216].

[9] K. Higashijima, T. Kimura and M. Nitta, Nucl. Phys. B **645**, 438 (2002) [arXiv:hep-th/0202064].

[10] K. Higashijima, T. Kimura and M. Nitta, Nucl. Phys. Proc. Suppl. **117**, 867 (2003) [arXiv:hep-th/0210034].

[11] M. Nitta, arXiv:hep-th/0309004.

[12] E. Kiritsis, C. Kounnas and D. Lust, Int. J. Mod. Phys. A **9**, 1361 (1994) [arXiv:hep-th/9308124].

[13] E. Witten, Phys. Rev. D **44**, 314 (1991).

[14] K. Hori and A. Kapustin, JHEP **0108**, 045 (2001) [arXiv:hep-th/0104202]; JHEP **0211**, 038 (2002) [arXiv:hep-th/0203147].

[15] K. Higashijima and E. Itou, Prog. Theor. Phys. **108**, 737 (2002) [arXiv:hep-th/0205036].

[16] K. Higashijima and E. Itou, Prog. Theor. Phys. **109**, 751 (2003) [arXiv:hep-th/0302090].

[17] K. Higashijima and E. Itou, Prog. Theor. Phys. **110**, 563 (2003) [arXiv:hep-th/0304194].

[18] E. Itou, to appear in the proceedings of 12th International Conference on Supersymmetry and Unification of Fundamental Interactions (SUSY 04), Tsukuba, Japan, 17-23 Jun 2004, [arXiv:hep-th/0409220].

[19] K. Higashijima and M. Nitta, Prog. Theor. Phys. **105**, 243 (2001) [arXiv:hep-th/0006027]; K. Higashijima, E. Itou and M. Nitta, Prog. Theor. Phys. **108**, 185 (2002) [arXiv:hep-th/0203081].

[20] I. Bakas, contributed to Annual International Conference on Strings, Theory and Applications (Strings 2004), Paris, France, 28 Jun - Jul 2, 2004, [arXiv:hep-th/0410093].

[21] K. Higashijima and M. Nitta, Prog. Theor. Phys. **103**, 635 (2000) [arXiv:hep-th/9911139]; Prog. Theor. Phys. **103**, 833 (2000) [arXiv:hep-th/9911225]; in Proceedings of Confinement 2000 (World Scientific, 2001), Eds. H.Suganuma et al., pp. 279 - 286, [arXiv:hep-th/0006025]; in Proceedings of ICHEP2000 (World Scientific 2001), Eds. C.S.Lim et al., pp. 1368 - 1370, [arXiv:hep-th/0008240].

[22] K. Higashijima, T. Kimura, M. Nitta and M. Tsuzuki, Prog. Theor. Phys. **105**, 261 (2001) [arXiv:hep-th/0010272].

[23] N. Nakayama and K. Sugiyama arXiv:hep-th/0411143.